



3299 North Wellness Drive
Building C Suite 240
Holland, Michigan
49424

phone 616.738.4420
fax 616.738.4432

Discharge Instructions Head Injury

Name: _____

David W. Lowry, MD, PC
M. Adam Kremer, MD, PLC
Scott M. Tuinstra PA-C, MS
Jefferson Alferink, PA-C
Neurosurgery

Medications:

- Take any medications only as directed.
- If you experience constipation while taking a narcotic pain reliever, begin using an over-the-counter laxative.

Diet:

- Resume whatever diet you were on before your injury slowly as you tolerate it (e.g., unrestricted, diabetic, low sodium, etc.)

Activity:

- Avoid pushing/pulling and lifting over 15 lbs. for four weeks.
- Do not participate in sports or physical education activities for four weeks.
- Walking is a good exercise. Go for at least four short walks a day, even if inside your home.
- You may remove your Ted Hose stockings once you are consistently walking four times a day.
- Do not drive before two weeks, if you periodically feel disoriented or if you have had any pain medication within four hours. You may ride in a car as a passenger as much as you tolerate.

CALL 911 FOR ANY SUDDEN OR RAPID WORSENING.

If Stable Call Our Office For:

- Increased drowsiness
- Worsening irritability
- Worsening nausea or vomiting
- Increasing headache
- New or worsened neck stiffness
- Seizure activity
- Slurring or incoherent speech
- Clear or bloody watery drainage for the nose or ears
- New or worsened weakness in either leg or arm

Follow-up:

- Follow up with your primary care physician unless otherwise directed by neurosurgeon.